

Research Article

Effects of integrated use of grass strip and soil bund on soil properties, Southern Ethiopia

Solomon Umer¹, Abebayehu Aticho^{2*}, Endalikachew Kiss²

¹ Department of Natural Resource Management, Werabe University, Werabe, Ethiopia

² Department of Natural Resource Management College of Agriculture and Veterinary Medicine, Jimma University, Jimma, Ethiopia

*corresponding author address: j.aticho@yahoo.com

Received 5 October 2018, Accepted 6 November 2018

Abstract: Soil degradation is the major environmental and agricultural production problem in the highlands of Ethiopia. The objectives of the study were to examine farmers' use of grass strip and soil bund to minimize soil degradation, and effects of grass strip and soil bund on basic soil properties. Structured questionnaire survey and field observations were conducted to collect data on farmers' use of grass strips and soil bund to minimize soil erosion, improve soil fertility and crop productivity. Simultaneously, composite soil samples were collected from croplands treated with both grass strip and soil bund, and the adjacent lands treated with soil bund only. The statistical result indicates respondents sex, marital status, educational level, erosion degree, access to conservation information, contact with extension workers, and trainings were significantly ($P < 0.05$) and positively affect the use of grass strips while livestock and land holding sizes were significantly ($P < 0.05$) and negatively affect farmers use of grass strip. The interaction of land management practices (grass strip and soil bund) and slope positions was significantly ($P < 0.05$) affects soil clay fractions and moisture, the highest amounts was observed at lands managed with grass strip situate at lower slope position. Similarly, soil organic matter, total nitrogen, available phosphorus, and cation exchange capacity were significantly ($P < 0.05$) influenced by the combined effects of land management practices and slope positions, highest quantity of these properties were observed lands managed with a grass strip and soil bund in lower slope position. Hence, integrated use of soil bund with grass strip has a meaningful contribution for selected soil property improvement that enhances soil fertility and crop productivity.

Keywords: *grass strip, land management, soil conservation practice, soil degradation, soil erosion*

To cite this article: Umer, S., Aticho, A. and Kiss, E. 2019. Effects of integrated use of grass strip and soil bund on soil properties, Southern Ethiopia. *J. Degrad. Min. Land Manage.* 6(2): 1569-1578, DOI: 10.15243/jdmlm.2019.062.1569.

Introduction

Water erosion induced soil degradation is one of the most destructive and widespread phenomena and realized as the key issue affecting food security. Of the total areas affected by soil degradation global, water erosion occupies 56% (Oldeman et al., 1991) and affects 80% of agricultural lands (Angima et al., 2003). Global assessment of human-induced soil degradation (GLASOD) reported, about 36% of farmlands are degraded at rates of 5-6 million ha per annum

(Scherr, 1999). The problem is severe in developing countries (like Africa and Asia), most of the lands in Africa are vulnerable to soil and environmental degradation (Vlek et al., 2008). Erosion-induced soil degradation, aggravate the problems of low agricultural productivity, food insecurity and rural poverty in Ethiopia (Smith 2010). In the highlands of Ethiopia (i.e., above 1500 m.a.s.l) erosion remove over 1.5 billion tons of top fertile soils annually (Taddese, 2001). Undulated terrain conditions and erosive rainfalls in the highlands of Ethiopia coupled with

detrimental farming practices worsen the problems of land degradation and make the soils more fragile. However, Ethiopian highlands hold greatest agricultural potentials and settlements; it accommodates over 88% of the human and 77% of the livestock populations of the country (Teklu, 2005). Because of this, water erosion is a serious threat to the development (e.g., economic, social, environmental) of Ethiopia (Tefera and Sterk, 2010; Amsalu and de Graaff, 2007; Pender et al., 2001; Shiferaw and Holden, 1999).

Since 1970's in collaboration with international partners, Ethiopia has made substantial efforts to introduce soil and water conservation technologies to reduce the risks of soil erosion and associated problems (Kebede, 2014). In Ethiopia the main challenges of soil and water conservation include, focus on initial construction of physical structures (e.g., terraces, bund, etc.), implement uniform technologies for all conditions (e.g., agro-ecological), fail to integrate physical structures with biological practices, top-down extension approach, and provide little attention to indigenous knowledge (Mitiku et al., 2006; Bekele et al., 2018). Because of this bottleneck, soil and water conservation efforts are less successful in the nation. Lately, the government has realized the cause failures in soil and water conservation efforts and give due attention to integrate physical structures with biological practices, enhance community engagement at various stages, and promote untapped indigenous soil conservation practices. Accordingly, in 2007 Ministry of Agriculture and Rural Development has adopted the framework of WOCAT (World Overview of Conservation Approaches and Technologies) to implement Sustainable Land Management (SLM) Program. As WOCAT framework focus on locally tried and tested techniques, the public and private partners use desho grass (*P. pedicellatum*) across the escarpment in densely populated highland for sustainable land management program (Smith, 2010).

Desho grass belongs to the family of Poaceae and native to tropical countries (Welle et al., 2006; Smith, 2010). This grass has an extensive root system and produces high biomass per unit area (Ramirez et al., 2010). In Ethiopia, the grass is discovered in 1991 at Chenecha District Southern region (Welle et al., 2006), and well adopted in different agroecology (Leta et al., 2013). This grass had been used as grass strip to protect croplands from soil erosion and degradations in highlands of Ethiopia (Welle et al., 2006; Yakob et al., 2015), to rehabilitate degraded land (Smith 2010), and to improve grazing land and livestock feeding (Welle et al.,

2006; Danano, 2007). As compared to other grasses (e.g., elephant *P. purpureum*, vetiver *V. zizanioides*) used for land management practices in Ethiopia, desho grass is among the most desirable one to control erosion and rehabilitate degraded land (Yakob et al., 2015). Based on the grass's character (e.g., root system) and diverse roles (erosion control, land rehabilitation, feed for cattle), desho grass is probably widely distributed and used for land management practices in Ethiopian highlands and others countries with similar agroecology. However, farmer's use of desho grass as grass strip to protect cropland from erosion, rehabilitate degraded land and livestock feeding is uneven among farmers. Farmers' use of land management practices are affected by institutional, technological, and socioeconomic factors (Kebede, 2014). In the study area, very limited information is available on farmers' use of desho grass as grass strip, and the effects of desho grass strip and soil bund at different slopes category on soil properties. Therefore, this study aims to assess farmers' use of desho grass as grass strip to minimize soil erosion and degradation, and examine the effects of grass strip and soil bund on selected soil properties at different slopes positions.

Materials and Methods

Description of the study area

This study area is located near to Analemmo District town Fonka Hadiya Zone, Southern Nation Nationalities and Peoples Regional State of Ethiopia. It is situated between 7.54- 7.70 latitude and 37.89 - 38.06 longitudes. Analemmo is 210 km away from Addis Ababa, the capital city of Ethiopia, and bounded with Silte Zone in North, Lemo District in South, Shashogo District in East, and Misha District in West direction (Figure 1). The study area receives mean annual rainfall between 1001-1200 mm and temperature within the ranges of 15-20°C. The rainfall distribution is characterized as bimodal; the small rains from March to April and the main (heavy) rain from June to October (AWOARD, 2013 unpublished). Most parts of Southern Ethiopia is dominated by schists, gneisses, granites and slates which are exposed in areas where geological erosion has been great (FAO 1986). The dominant soils of southern Ethiopia including of the study areas are Vitric and Nitisol (FAO/UNDP, 1984), Luvisols, Regosols and Cambisols (Adbacho, 1991). The average rate of population growth in South Ethiopia is 2.9% per annum with an average population density of 127 per sq.km. While in Hadiya Zone, the population density is

346 per sq.km, which is to be overpopulated and higher than the regional average (SNNPRSO, 2006). Mixed agriculture (integration of crop and livestock) is livelihood base of the community.

Methodological approach

To understand the effects of desho grass used as grass strip to stabilize soil bund at different slope positions on soil properties, composite soil samples were collected from croplands situated at upper and lower slope positions treated with both grass strip and soil bund, and adjacent cropland managed with soil bund alone (Figure 2). The sample plots were used for wheat production with uniform agronomic practices such as nutrient management and weed control practices. As presented in Table 1, total 16 composite soil samples were collected 30 cm sampling depths using an auger for chemical property and nutrient content analysis. At the same time, 16 undisturbed soil samples were collected for bulk density and moisture content determination. Past studies on farmer's participation and use of land management practices, provide lists of variables that influence farmers decision (Ervin, 1982; Tadesse and Belay, 2005). Based on information obtained from literatures, in this study, the following conceptual framework was adopted to capture variables influence grass strip use for land management practices in the study area (Figure 3).

A structured questionnaire survey was used to identify factors affecting farmers (households) use of grass strip as land management options. A simple random sampling technique was employed to choose respondent households for a face-to-face interview.

Table 1. Description of soil samples collected for laboratory analysis

| Land management practices | Slope position | Replication |
|-----------------------------|----------------|-------------|
| Soil bund alone | Upper | 4 |
| | Lower | 4 |
| Grass strip and soil bund | Upper | 4 |
| | Lower | 4 |
| Total composite soil sample | | 16 |

The farmer's list existing in village administrative centre (locally called Kebele-the smallest administrative unit of Ethiopia) has used as a sample frame for random sampling techniques. The respondent sample size was determined using Cochran (1977) formula provided for finite population (below 10000). Accordingly, 110 randomly selected respondent households were interviewed.




Figure 1. Geographical map of the study area



Figure 2. Partial views of bund stabilized with grass strip and bund alone




Figure 3. Conceptual framework

Soil laboratory analysis

The soil samples collected from the study area were transported to Jimma University College of Agriculture and Veterinary Medicine research and teaching soil laboratory. Before analyzing soil texture, pH, EC, CEC, OC, TN, and Av. P samples were air dried, grounded, and sieved with standard mesh size. Soil particle size distribution was determined by Boycouse Hydrometer method (Houba et al., 1989) after organic matter was destroyed by addition of hydrogen peroxide (H₂O₂), disperse cementation with hexametaphosphate (NaPO₃)₆ and sodium carbonate (Na₂CO₃), and removes foam via amyl alcohol. Bulk density of undisturbed soil sample were determined by core method (FAO, 2007), using the ratio of solid mass to total volume when the sample was dried in an oven at 105°C for 24 hours. The gravimetric method was used to determine soil moisture content. Soil reaction (pH) was measured with pH meter glass electrodes in 1:2.5 (soil: water on a mass to

volume basis) suspension after calibration of the pH meter with buffer solutions (pH 4, 7 and 10). Soil electrical conductivity (EC) was measured by conductivity meter 1:2.5 (soil: water on a mass to volume basis) suspension after the calibration of EC meter. Walkley and Black wet digestion method was used for organic carbon determination (Walkley and Black, 1934). Soil total nitrogen was determined by Kjeldahl digestion procedure (Bremmer, 1996). Bray II method was used to extract available phosphorus (Av. P) from soil (Van Reeuwisk, 1993), and the extracted Av. P was determined using spectrophotometer at a wavelength of 820 nm. While ammonium acetate method (1N NH₄OAc) at pH 7.0 was used to determine cation exchange capacity (CEC) of the soil (Houba et al., 1989).

Statistical analysis

Both laboratory and survey data were subjected SPSS (Statistical Package for Social Science, version 20 software). Soil laboratory data was

statistically analyzed via two ways ANOVA, the main and interaction effects of land management practices (grass strip with soil bund, and soil bund alone) and slope positions (upper and lower) on soil properties, and significant means ($P = 0.05$) were separated using LSD. The t-test and chi-square were used to analyze continuous variables, and dummy variables gathered via survey, respectively.

Results and Discussion

Farmers use of grass strip to stabilize soil bunds

In the study area, soil bund was widely constructed on croplands to collect surface runoff, increase infiltration, and minimize soil erosion. About 70.9% of the respondent households use desho grass as grass strip on soil bunds while the rest 29.1% left the bund bare. The statistical result indicates household land and livestock holding sizes were significantly ($P < 0.05$) and negatively influence the use of desho grass strip to stabilize bunds (Table 2). As land and livestock holding size of the household gets larger, the probability to use grass strips to minimize erosion on

croplands becomes lower. Perhaps this could be, farmers possessed large cropland requires huge bunches (bundles) of grass to establish grass strips on their croplands - which requires huge amount of money to buy grass. On the other hand, large livestock sizes can destroy the grass strip while grazing on the croplands after harvest. Because of this, smaller land and livestock holders use grass strips to stabilize soil bunds, minimize erosion and enhance soil productivity. This argues with the finding of Sinore et al. (2018), who stated that land and livestock holding size has a strong association with farmer's perception on the uses of biological land management practices such as grass strips to minimize land degradation problems and enhances land productivity. This disagreement is arising due to the use of similar land management practice for lands under different conditions: already degraded land for restoration and stable cropland for maintaining productivity. Accordingly, the current study deal with the use of grass strips (land management) to preserve stabile croplands for sustainable production, while the former study deal with the use of land management practices to restore degraded lands.

Table 2. Independent t-test on factors influence grass strip to stabilization soil bund

| Dependant variables | | Mean | S.D | t-test | Sig. |
|---------------------------|------------------|-------|-------|--------|-------|
| Livestock holding (TLU) | Grass strip user | 2.97 | 1.02 | -2.56 | 0.012 |
| | Non- user | 3.62 | 1.58 | | |
| Land holding size (ha) | Grass strip user | 0.71 | 0.38 | -3.39 | 0.001 |
| | Non- user | 1.03 | 0.58 | | |
| Farming experience (year) | Grass strip user | 29.74 | 11.07 | 1.40 | 0.165 |
| | Non- user | 26.53 | 10.58 | | |
| Family size (number) | Grass strip user | 1.67 | 0.77 | 0.12 | 0.900 |
| | Non- user | 1.69 | 0.86 | | |

Furthermore, household head sex, marital status, educational level, cropland erosion status, information access, contact with extension workers, and trainings were significantly ($P < 0.05$) influence farmers use of grass strip for land management purpose (Table 3). Unlike married farmers, the unmarried have poor interest to invest in long-term land management practices (e.g. grass strip) and focus on short-term gains. In the society, married farmers carry more social, economic and environmental responsibility than the unmarried farmers did. Oakley and Garforth (1997) reported unmarried men in some communities are expected to work on his parent's farm, and only married are expected to start farming his own to feed his family members. The educated households tend to use grass strips to prevent soil erosion and improve soil productivity.

Because the length of attending formal education enables farmers to easily realize extensions information and apply land management options to enhance crop production and productivity to realize household food security and generate sufficient income. This argues with the finding of Budry et al. (2006) who reported, more educated farmers place greater emphasis on finding jobs outside the community instead of managing soil conservation structures on their plots. As compared to Western construes, in Ethiopia getting a better off-farm job (e.g., government, non-government, and private sectors) is very difficult for farmers who graduated high school. Thus, most high school graduates mainly depend on agriculture to produce food for home consumption and generate incomes through properly managing lands.

Table 3. Chi-square test on the relationship between grass strip user and non-user farmers

| Variables | | User | Non-user | P-value |
|-------------------------|--------------------------|-------|----------|---------|
| Sex | Male | 61.8% | 10.9% | 0.000 |
| | Female | 9.1% | 18.2% | |
| Age | Productive (> 18 < 65) | 40.0% | 13.6% | 0.362 |
| | Unproductive (< 18 > 65) | 30.9% | 15.5% | |
| Marital status | Married | 61.8% | 21.8% | 0.019 |
| | Single | 0.9% | 4.5% | |
| | Divorced | 2.7% | 1.8% | |
| | Widowed | 5.5% | 0.9% | |
| Educational status | Illiterate | 5.5% | 2.7% | 0.012 |
| | Primary | 10.9% | 11.8% | |
| | Secondary | 54.5% | 14.5% | |
| Cropland erosion degree | Sever | 68.2% | 2.7% | 0.0001 |
| | Moderate | 2.7% | 26.4% | |
| SWC information access | Development agents (DA) | 47.3% | 1.8% | 0.001 |
| | Family in neighbour | 3.6% | 8.2% | |
| | Media (FM radio) | 13.6% | 13.6% | |
| | NGOs | 6.4% | 5.5% | |
| Contact with DA | Once a week | 14.5% | 8.2% | 0.051 |
| | Twice a week | 15.5% | 11.8% | |
| | Once a month | 40.9% | 9.0% | |
| Attend SWC training | Yes | 70.0% | 0.9% | 0.001 |
| | No | 0.9% | 28.2% | |

In the study area, farmers' access to extension information (e.g., media, training, extension workers) improves farmers understanding of land management options and facilitates their land management decisions. This coincides with previous studies, access to agricultural information improves farmers adoption soil conservation (Yishak, 2005; Batiwaritu and Mvena, 2009).

Effects of grass strip on selected soil physical properties

Statistical analysis of main effects revealed, clay fractions were soil significantly ($P < 0.01$) differ as function of land management practices (grass strip and soil bund, and soil bund alone) and slope positions, more amount was found at soil bunds stabilized with grass strip situated at lower slope positions (Table 4). Similarly, the interaction effects of land management practices and slope positions significantly ($P < 0.05$) influences clay fractions, the highest amount of clay was recorded at soil bunds stabilized with grass strips situated in lower slope positions. The low clay fraction at bunds treated with grass strip in the upper slope (42.0 ± 2.58) position was not an indicator the poor performance of grass strip to capture clay materials in the upper slope position. However, this confirms the effects of past water erosion events or topographic effects on clay fractions. The past erosion events caused the removal of

clay fraction from the upper slope position, are transported and deposited into lower slope positions; consequently, more sand material is remained in upper slope positions. It was evidenced that at the upper slope positions soil bunds managed with grass strip had relatively better clay (42.0 ± 2.58) than soil bund alone (41.0 ± 2.23). This indicates that using grass strip as land management option at the upper and lower slope position could have similar soil protection effects. Other studies reported soil bunds stabilized with grass are contributes for high proportion of clay particles than the bund alone (Tadele, 2011) and fine materials are selectively removed from the upper slope and deposited at the lower slope (Tripathi and Singh, 2001). The main effect of both land management practices and slope position significantly ($P < 0.05$) contributed for soil moisture and bulk density improvement. However, the interaction effect shows land management practices and slope position significantly ($P < 0.01$) influences soil moisture. The highest soil moisture was recorded from soil bunds managed with grass strip situated at lower slope position (18.04 ± 0.54) while the lowest at upper slope position (15.13 ± 0.12). This could be due to the presence of high clay fraction in lower slope positions. Furthermore, root systems of the grass strips conserve moisture through minimizing erosion, evaporation, surface runoff and modifying soil microenvironment.

Table 4. Soil physical properties as influenced by integrated use grass strip and soil bunds at different slope position

| Soil physical property | Mean ± SD | P-value | Interaction of LMP x SP | Mean ± SD | P-value |
|-------------------------------|-------------------------|---------|-------------------------------------|---------------------------|---------|
| LMP | | | | | |
| Sand (%) SBG | 20.75±3.19 ^b | 0.0001 | Sand (%) SBG x upper | 22.15±1.9 ^b | 0.012 |
| Sand (%) SB | 22.00±2.82 ^a | | Sand (%) SBG x lower | 18.5± 1.09 ^c | |
| Silt (%) SBG | 33.25±2.76 ^a | 0.4421 | Sand (%) SB x upper | 23.5± 2.51 ^a | |
| Silt (%) SB | 34.25±2.81 ^a | | Sand (%) SB x lower | 21±3.41 ^{ab} | |
| Clay (%) SBG | 44±3.85 ^a | 0.0005 | Silt (%) SBG x upper | 34.5±2.51 ^a | 0.337 |
| Clay (%) SB | 43 ±4.11 ^b | | Silt (%) SBG x lower | 36.05±3.41 ^a | |
| BD (g/cm ³) SBG | 1.06±0.05 ^b | 0.0006 | Silt (%) SB x upper | 33.15± 3.40 ^a | |
| BD (g/cm ³) SB | 1.11±0.03 ^a | | Silt (%) SB x lower | 35.25±2.51 ^a | |
| MC (%) SBG | 16.02±1.43 ^a | 0.0002 | Clay (%) SBG x upper | 42.0±2.58 ^c | 0.003 |
| MC (%) SB | 12.10±1.54 ^b | | Clay (%) SBG x lower | 49.0± 1.63 ^a | |
| Slope position (SP) | | | | | |
| Sand (%) upper | 23.00±2.13 ^a | 0.0004 | Clay (%) SB x upper | 41.0± 2.23 ^{ab} | |
| Sand (%) lower | 19.75±2.91 ^b | | Clay (%) SB x lower | 45.0± 1.33 ^b | |
| Silt (%) upper | 32.00±2.77 ^b | 0.0813 | BD (g/cm ³) SBG x upper | 1.05±0.05 ^a | 0.061 |
| Silt (%) lower | 33.00±2.13 ^b | | BD (g/cm ³) SBG x lower | 1.03± 0.03 ^a | |
| Clay (%) upper | 45.00±2.13 ^b | 0.0003 | BD (g/cm ³) SB x upper | 1.10± 0.016 ^a | |
| Clay (%) lower | 47.75±3.37 ^a | | BD (g/cm ³) SB x lower | 1.11±0.08 ^a | |
| BD (g/cm ³) upper | 1.15±0.04 ^a | 0.0005 | MC (%) SBG x upper | 15.13± 0.12 ^b | 0.001 |
| BD (g/cm ³) lower | 1.07±0.05 ^b | | MC (%) SBG x lower | 18.04±0.54 ^a | |
| MC (%) upper | 10.53±2.56 ^b | 0.0002 | MC (%) SB x upper | 12.09± 0.82 ^{ab} | |
| MC (%) lower | 11.58±1.65 ^a | | MC (%) SB x lower | 14.17±0.46 ^c | |

LMP = land management practices; SBG = soil bund integrated with grass strip, SB= soil bund alone, in column two superscript followed with a same lower letter for the same soil parameters are not significantly different at P=0.05

This agrees with Danano (2007) who reported, the root system of desho grass holds soil particles, improves structure, and soil water retention.

Effect of grass strips on soil chemical properties

Statistical analysis shows main effects of land management practices (grass strip and soil bund, soil bund alone) and slope positions significantly (P< 0.05) influence soil pH, OC, N, C:N, Av. P, and CEC (Table 5). However, interaction effects of land management practices and slope positions significantly (P < 0.05) influence soil OC, TN, C:N, Av. P, and CEC. The highest amounts of soil OC, TN, C:N, Av. P, and CEC were observed at soil bunds managed with grass strips in lower slope position. This implies using grass strip as alternative land management options has a meaningful contribution for soil nutrient (e.g., Av. P), soil organic matter content (via above and below ground biomass addition), nutrient retention capacity (CEC), and organic matter decomposition. Other findings also witnessed

about desho grass contribution for soil chemical property improvement through its extensive root system that anchors soil particles, high biomass production capacity and return organic matter to soil (Ramirez et al., 2010), minimize and control soil and nutrient loss (Welle et al., 2006; Yakob et al., 2015), and rehabilitate degraded area (Smith 2010). Moreover, other authors reported the presence of high nutrient in the accumulation zone (lower slope position) than the loss zone (upper slope position) (Kibret, 2008; Tadele et al., 2011). The highest amounts of OC, TN, C:N, Av. P, and CEC recorded in the lower slope position were not only due to the interaction effects of land management practices and slope position but also from past deposition events. For instance, when the quantity of soil properties (e.g., Av. P, C:N, CEC.) observed in soil bunds managed with a grass strip, and soil bunds alone at the upper slope positions, a better property was found at soil bunds managed with a grass strip.

Table 5. Effects of desho grass integration with bunds on at different slope position soil physical properties

| Soil chemical property | Mean ± SD | P-value | Interaction (LMP x SP) | Mean ± SD | P-value |
|------------------------------|--------------|---------|---------------------------------|--------------|---------|
| LMP | | | | | |
| pH-H ₂ O SBG | 5.69±0.26a | 0.0001 | pH-H ₂ O SBG x upper | 6.00± 0.00a | 0.2732 |
| pH-H ₂ O SB | 5.36±0.08b | | pH-H ₂ O SBG x Lower | 5.50± 0.54a | |
| EC (ds/m) SBG | 0.128±0.03a | 0.0623 | pH-H ₂ O SB x upper | 5.00± 0.05a | |
| EC (ds/m) SB | 0.137±0.02a | | pH-H ₂ O SB x lower | 5.44± 0.16a | |
| OC (%)SBG | 2.33±0.44a | 0.0010 | EC (ds/m) SBG x upper | 0.055±0.00a | 0.1010 |
| OC (%)SB | 1.85±0.37b | | EC (ds/m) SBG x Lower | 0.115±0.02a | |
| TN (%)SBG | 0.20±0.03a | 0.0001 | EC (ds/m) SB x upper | 0.082±0.12a | |
| TN (%)SB | 0.19±0.03b | | EC (ds/m) SB x lower | 0.142±0.02 a | |
| C: N SBG | 11.65±0.02a | 0.001 | OC (%) SBG x upper | 1.51±0.98ab | 0.0001 |
| C: N SB | 9.73±0.01b | | OC (%) SBG x Lower | 2.48± 0.15a | |
| Av.P (ppm) SBG | 6.12±0.78a | 0.0001 | OC (%) SB x upper | 2.18±0.10b | |
| Av.P (ppm) SB | 4.84±0.81b | | OC (%) SB x lower | 1.69± 0.13c | |
| CEC(cmol+)/kg SBG | 28.82±2.99a | 0.0218 | TN (%) SBG x upper | 0.13±0.017ab | 0.0310 |
| CEC(cmol+)/kg SB | 26.92±2.17b | | TN (%) SBG x Lower | 0.22±0.01a | |
| Slope position(SP) | | | TN (%) SB x upper | 0.18± 0.016b | |
| pH-H ₂ O at upper | 5.63±0.31a | 0.0300 | TN (%) SB x lower | 0.16±0.02c | |
| pH-H ₂ O at lower | 5.41±0.11b | | C:N SBG x upper | 15.75±0.021b | 0.0001 |
| EC (ds/m) at upper | 0.098±0.054a | 0.9336 | C:N SBG x Lower | 17.2±0.021a | |
| EC (ds/m) at lower | 0.097±0.028a | | C:N SB x upper | 12.11±0.062c | |
| OC (%) at upper | 1.60±0.144b | 0.0001 | C:N SB x lower | 10.56±0.06ab | |
| OC (%) at lower | 2.00±0.202a | | Av.P(ppm) SBG x upper | 5.00± 0.00b | 0.0001 |
| TN (%) at upper | 0.14±0.026b | 0.0001 | Av.P(ppm) SBG x Lower | 6.50± 0.57a | |
| TN (%) at lower | 0.16±0.02a | | Av.P (ppm) SB x upper | 4.00± 0.00ab | |
| C:N at upper | 11.65±0.02a | 0.001 | Av.P (ppm) SB x lower | 4.09± 0.14c | |
| C:N at lower | 9.73±0.01b | | CEC(cmol+)/kg SB x upper | 30.00±2.30b | 0.0048 |
| Av.P (ppm) upper | 6.12±0.78a | 0.0001 | CEC(cmol+)/kg SBG x lower | 31.50±1.29a | |
| Av.P (ppm) lower | 4.84±0.81b | | CEC(cmol+)/kg SB x upper | 25.75±0.95ab | |
| CEC(cmol+)/kg upper | 26.35±1.40b | 0.0009 | CEC(cmol+)/kg SB x lower | 27.25±0.95c | |
| CEC(cmol+)/kg lower | 30.39±1.74a | | | | |

LMP = land management practices; SBG = soil bund integrated with a grass strip, SB= soil bund alone, in column two superscript followed with a same lower letter for the same soil parameters are not significantly different at P=0.05

Conclusions

Integrating soil bund with grass strip as land management options play important roles in maintaining land productivity or restore degraded lands situated at different slope position. It has improved soil properties (like CEC, OC, Av. P) at croplands found in upper and lower slope positions. The difference observed in soil physical and chemical properties between slope positions (lower and lower) is not due to grass strip malfunction at slope positions but caused by past erosion events and topographic effects on soil property. Using desho grass strip could have relatively similar effects on soil properties at different slope positions. However, farmers' socio-economic status, cropland erosion, and access to information are affecting the use of desho grass strip to ensure sustainable land management practices. General, to scale-up desho grass strip as one of land management options in

similar agro-ecological setting desho grass strip specific extension has to be provided to the community.

Acknowledgement

The authors would like to thank Jimma University College of Agriculture and Veterinary Medicine for financial support of the research work.

References

- Adbacho, W.1991. Biological or Vegetation Cover as a Sustainable Soil Conservation Strategy: A case study of Tilku Demala, M.A Thesis: School of Environmental Science; University of East Anglia.
- Amsalu, A. and de Graaff, J. 2007. Determinants of adoption and continued use of stone terraces for soil and water conservation in an Ethiopian highland watershed. *Ecological Economics* 61:294–302.
- Angima, S.D., Stott, D.E., O'Neill, M., Ong, C.K. and Weesies, G.A. 2003. Soil erosion prediction using

- RUSLE for central Kenya highland conditions. *Agriculture, Ecosystems and Environment* 97: 295–308 doi:10.1016/S0167-8809(03)00011-2.
- Batiwaritu, G. and Mvena, Z.S. 2009. Determinants of the adoption of physical soil bund conservation structures in Adama District, Oromia Region, Ethiopia. *East Africa Journal of Sciences* 3:142-152.
- Bekele, A., Aticho, A. and Kissi, K. 2018. Assessment of community based watershed management practices: emphasis on technical fitness of physical structures and its effect on soil properties in Lemo district, Southern Ethiopia. *Environmental Systems Research* 7:20, doi: <https://doi.org/10.1186/s40068-018-0124-y>.
- Bremner, J.M. 1996. Nitrogen Total. In: Sparks, D.L. (ed), *Methods of Soil Analysis Part 3: Chemical Methods*, SSSA Book Series 5, Soil Science Society of America, Madison, Wisconsin, 1085-1122.
- Budry, B., Curtis, M. and Dennis, A.S. 2006. The adoption and management of soil conservation practices in Haiti: the case of rock walls. *Agricultural Economics Review* 7(2) http://www.eng.auth.gr/mattas/7_2.htm.
- Cochran, W.G. 1977. *Sampling technique*. 3rd ed. New York, John Wiley Sons. p 442.
- Danano, D. 2007. Improved grazing land management-Ethiopia. In: Liniger, H. and Critchley, W. (eds), *Where the land is greener*. Bern, Switzerland: WOCAT. pp. 313–316.
- Ervin, D.E. 1982. Soil erosion control on owner-operated and rented cropland. *Journal of Soil and Water Conservation* 37: 285-288.
- FAO. 1986. Ethiopian highlands reclamation study. Final report prepared for the Government of Ethiopia Volume 1. Rome, Italy. p 354.
- FAO. 2007. *Methods of Analysis for Soils of Arid and Semi-Arid Regions*. Food and Agriculture Organization of the United Nations Rome, Italy.
- FAO/UNDP. 1984. Ethiopian geomorphology and soils. Food and Agriculture Organization of the United Nations/United Nations Development Program Assistance to Land Use Planning. Addis Ababa. p 161.
- Houba, V.J.G., Van der Lee J.J., Novozamsky, I. and Walinga, I. 1989. Soil and plant analysis, a series of syllabi, part 5; soil analysis procedures. Wageningen Agricultural University, Wageningen, The Netherlands.
- Kebede, W. 2014. Effect of soil and water conservation measures and challenges for its adoption: Ethiopia in focus. *Journal of Environmental Science and Technology* 7: 185-199. DOI:10.3923/jest.2014.185.199.
- Kibret, M. 2008. Enclosure as a Viable Option for Rehabilitation of Degraded Lands and Biodiversity Conservation: The Case of Kallu Woreda, Southern Wello. M.Sc. Thesis Addis Ababa University.
- Leta, G., Duncan, A. and Abdena, A. 2013. Desho grass (*Pennisetum pedicellatum*) for livestock feed, grazing land and soil and water management on small-scale farms. NBDC Brief 11. Nairobi, Kenya: ILRI.
- Mitiku, H., Herweg, K. and Stillhardt, B. 2006. Sustainable land management: a new approach to soil and water conservation in Ethiopia. Mekelle, Ethiopia: land resources management and environmental protection department, Mekelle University; Bern, Switzerland: center for development and environment, University of Bern, and Swiss national center of competence in research (NCCR) North South. p 269.
- Oakley, P. and Garforth, C. 1997. Agriculture Extension and Rural Development Centre, School of Education, University of Reading, UK. Food and Agriculture Organization of the United Nations Rome, Italy.
- Oldeman, L.R., Hakkeling, R.T.A. and Sombrou, W.G. 1991. World Map of the Status of Human-Induced Soil Degradation: An Explanatory Note, UNEP and ISRIC, Wageningen. The Netherlands.
- Pender, J.L., Gebremehedhin, B., Benin S. and Ehui, S. 2001. Strategies for sustainable development in Ethiopian highlands. *American Journal of Agricultural Economics* 83: 1231–1240.
- Ramirez, K.S., Lauber, C.L., Knight, R., Bradford, M.A. and Fierer, N. 2010 Consistent effects of nitrogen fertilization on soil bacterial communities in contrasting systems. *Ecology* 91: 3463–3470.
- Scherr, S.J. 1999. Soil degradation, a threat to developing-country food security by 2020? Food, agriculture and the environment discussion paper 27. International Food Policy Research Institute 2033 K Street, N.W. Washington DC 20006- 1002 USA, pp 13-45.
- Shiferaw, B. and Holden, S.T. 1999. Soil erosion and smallholders' conservation decisions in the highlands of Ethiopia. *World Development* 27: 739–752.
- Sinore, T., Kissi K. and Aticho, A. 2018. The effects of biological soil conservation practices and community perception toward these practices in the Lemo District of Southern Ethiopia. *International Soil and Water Conservation Research* 6: 123–130.
- Smith, G. 2010. Ethiopia: local solutions to a global problem. <http://www.new-ag.info/en/focus/focusItem.php?a=1784>. (Accessed 4/11/2017 9:17 AM)
- SNNPRSO. 2006. Regional Atlas. Southern Nation, Nationalities and Peoples Regional State, Bureau of Finance and Economic Development, Bureau of Statistics and population. Awassa, Ethiopia.
- Tadele, A., Yihenuw, G.S., Mitiku, H. and Yamoh, C. 2011. Effect of soil and water conservation measures on selected soil physical and chemical properties and barley (*Hordeum spp.*). *Yield* 5: 1483-1495.
- Tadesse, M. and Belay, K. 2005. Factors influencing adoption of soil conservation measures in southern Ethiopia the case of Ginuno area. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*. 105: 49-62.
- Tefera, B. and Sterk, G. 2010. Land management, erosion problems and soil and water conservation in Fincha'a watershed, western Ethiopia. *Land Use Policy* 27: 1027–1037.

- Teklu, E. 2005. Land preparation methods and soil quality of a vertisol area in the central highlands of Ethiopia. PhD thesis, University of Hohenheim, Stuttgart.
- Tripathi, S.K. and Singh, K.P. 2001. Ecological responses of dry tropical forest and savanna ecosystems to nutrient enrichment. In: Jha, P.K., Baral, S.K., Karmacharya, S.B., Lekhak, H.D., Locoul, P. and C.B. Baniya, C.B.(eds), *Environment and Agriculture: Biodiversity, Agriculture and Pollution in South Asia*. Published by Ecological Society of Nepal, pp. 150-157.
- Van Reeuwijk, L.P. 1993. Procedures for Soil Analysis. 3rd Edition. International Soil Reference and Information Centre Wageningen (ISRIC). Wageningen The Netherlands.
- Vlek, P.L.G., Le, B.Q. and Tamene, L. 2008. Land decline in land-rich Africa. A creeping disaster in the making. Center for development research (ZEF) University of Bonn, Science Council and CGIAR, pp 12-54.
- Walkley, A. and Black, C.A. 1934. An examination of Degtjareff methods for determining soil organic matter and the proposed modifications of the chromic acid titration method. *Soil Science* 37:29-38.
- Welle, S., Chantawarangul, K., Nontananandh, S. and Jantawat, S. 2006. Effectiveness of grass strips as barriers against runoff and soil loss in Jijiga area, northern part of Somalia region, Ethiopia. *Kasetsart Journal: Natural Science* 40:549-558.
- Yakob, G., Gebremicheal, A., Aklilu, A. and Melaku, E. 2015. Participatory evaluation of different multipurpose grass species for graded soil bund stabilization in Gimbo District, South West Ethiopia. *Open Access Library Journal* 2:e1627. <http://dx.doi.org/10.4236/oalib.1101627>.
- Yishak, G. 2005. Determinants of Adoption of Improved Maize Technology in Damot Gale Woreda, Wolaita, Ethiopia, MSc Thesis submitted to Alemaya University.